

Groupschemes

Christian Dahlhausen

Transcript of

Christian Merten (cmerten@mathi.uni-heidelberg.de)

WiSe 2023

Contents

1	Introduction	3
----------	---------------------	----------

Chapter 1

Introduction

Literature: Görtz-Wedhorn: Algebraic Geometry I and II

The goal of this lecture is a brief introduction to the theory of group schemes.

Definition 1.1 (Group object). Let $\mathcal{C}$ be a category with finite products. A *group object in $\mathcal{C}$* is the data (G, m, e, i) where

- G is an object of $\mathcal{C}$
- $m: G \times G \rightarrow G$ is the multiplication map
- $e: 1 \rightarrow G$ is the unit
- $i: G \rightarrow G$ is the inversion map

such that the following diagrams commute

$$\begin{array}{ccc} G \times G \times G & \xrightarrow{m \times \text{id}} & G \times G \\ \downarrow \text{id} \times m & & \downarrow m \\ G \times G & \xrightarrow{m} & G \end{array}, \quad \begin{array}{ccc} G \times G & \xrightarrow{m} & G \\ \uparrow \text{id} \times e & \swarrow & \\ G \times 1 & & \end{array} \quad \text{and} \quad \begin{array}{ccc} G & \xrightarrow{\text{id} \times i} & G \times G \\ \downarrow & & \downarrow m \\ 1 & \xrightarrow{e} & G \end{array}.$$

G is called *commutative*, if additionally the diagram

$$\begin{array}{ccc} G \times G & \xrightarrow{\text{swap}} & G \times G \\ \downarrow m & \swarrow m & \\ G & & \end{array}$$

commutes.

A *morphism of group objects* $(G, m, e, i) \rightarrow (G', m', e', i')$ is a morphism $f: G \rightarrow G'$ in $\mathcal{C}$ such that the diagrams

$$\begin{array}{ccc} G \times G & \xrightarrow{f \times f} & G' \times G' \\ \downarrow m & & \downarrow m' \\ G & \xrightarrow{f} & G' \end{array}, \quad \begin{array}{ccc} G & \xrightarrow{f} & G' \\ \uparrow e & \swarrow e' & \\ 1 & & \end{array} \quad \text{and} \quad \begin{array}{ccc} G & \xrightarrow{f} & G' \\ \downarrow i & & \downarrow i' \\ G & \xrightarrow{f} & G' \end{array}.$$

This defines the category $\text{Grp}(\mathcal{C})$ of group objects in $\mathcal{C}$.

Example 1.2. • $\text{Grp}(\text{Set}) \simeq \text{Grp}$

- $\text{Grp}(\text{Grp}) \simeq \text{Ab}$
- $\text{Grp}(\text{Ab}) \simeq ?$

- $\text{Grp}(\text{Top}) \simeq$ topological Groups
- $\text{Grp}(\text{Mfd}_\infty) \simeq$ Lie Groups

Definition 1.3 (group scheme). Let S be a scheme. An S -group scheme or an S -group is a group object in the category of schemes over S .

Remark 1.4. Let S be a scheme. The structure of a group scheme over S on a S -scheme G is equivalent to a factorisation of the functor of points

$$\begin{array}{ccc} \text{Sch}_S & \longrightarrow & \text{Set} \\ \downarrow & \nearrow & \\ \text{Grp} & & \end{array}$$

via the forgetful functor from groups to sets.

Example 1.5. Let S be a scheme.

- (i) Let Γ be a group. Then $G = \Gamma_S$ where $G(T) := \{ \text{locally constant maps } T \rightarrow \Gamma \}$
- (ii) (additive group) $\mathbb{G}_{a,S}$ where $\mathbb{G}_{a,S}(T) = \mathcal{O}_T(T)$. We have $\mathbb{G}_{a,S} \simeq \mathbb{A}_S^1$.
- (iii) (multiplicative group) $\mathbb{G}_{m,S}$ where $\mathbb{G}_{m,S}(T) := \mathcal{O}_T(T)^\times$.
- (iv) (roots of unity) $\mu_{n,S}$ ($n \geq 1$) where $\mu_{n,S}(T) = \{x \in \mathcal{O}_T(T)^\times \mid x^n = 1\}$.
- (v) $S = \text{Spec}(R)$. $\text{GL}_{n,R} = \text{Spec}(A)$ where $A = R[T_{ij} \mid 1 \leq i, j \leq n][\det^{-1}]$ where $\det = \sum_{\sigma \in S_n} \text{sgn}(\sigma) T_{1\sigma(1)} \cdots T_{n\sigma(n)}$. We obtain $\text{GL}_{n,S}$ by base changing $\text{GL}_{n,\mathbb{Z}}$.

Lemma 1.6. Let G be a S -group. Then $G \rightarrow S$ is separated if and only if $S \xrightarrow{e} G$ is a closed immersion.

Definition 1.7. Let R be a ring. A (commutative) Hopf-Algebra over R is a group object in Alg_R^{op} , where $\text{Alg}_R = \text{CRing}_R$.

Remark 1.8. For a R -Hopf-Algebra A , we denote the canonical maps by

- $\mu: A \rightarrow A \otimes_R A$ (Comultiplication)
- $\varepsilon: A \rightarrow R$ (Counit)
- $\iota: A \rightarrow A$ (Antipode)

A Hopf-Algebra is called *cocommutative*, if the associated group object in Alg_R^{op} kommutativ ist.

Remark 1.9. For a ring R , by construction we have an equivalence of categories between the category of affine R -group schemes and the opposite category of R -Hopf-Algebras.

Example 1.10. The additive group $\mathbb{G}_{a,R} = \text{Spec}(R[t])$ has

- comultiplication $\mu: R[t] \rightarrow R[t] \otimes_R R[t], t \mapsto t \otimes 1 + 1 \otimes t$.
- counit $\varepsilon: R[t] \rightarrow R, t \mapsto 0$
- antipode $\iota: R[t] \rightarrow R[t], t \mapsto -t$

Proof. For any R -algebra A we have $\mathbb{G}_{a,R}(A) = A$ and the diagram

$$\begin{array}{ccc} \mathbb{G}_{a,R}(A) \times \mathbb{G}_{a,R}(A) & \xrightarrow{m} & \mathbb{G}_{a,R}(A) \\ \downarrow \simeq & & \downarrow \simeq \\ \text{Hom}_R(R[s_1, s_2], A) & \xrightarrow{\mu^*} & \text{Hom}_R(R[t], A) \end{array} .$$

□

Definition 1.11. Let G be a S -group. A *subgroupscheme* of G is a subscheme $H \subseteq G$ such that

- 1) for all $T \in \text{Sch}_S$, we have $H(T) \subseteq G(T)$ a subgroup,
- 2) We have commutative diagrams

$$\begin{array}{ccc} H \times_S H & \longrightarrow & G \times_S G \xrightarrow{m} G \\ \downarrow & \nearrow & \downarrow \\ H & & H \end{array} \quad \text{and} \quad \begin{array}{ccc} S & \xrightarrow{e} & G \\ \downarrow & \nearrow & \downarrow \\ H & & H \end{array}$$

A subgroup scheme $H \subseteq G$ is *normal* if $H(T)$ is a normal subgroup of $G(T)$ for all $T \in \text{Sch}_S$.

For a morphism $f: G \rightarrow G'$ of S -groups and a subgroup $H' \subseteq G'$, let $f^{-1}(H')$ be $G \times'_{G'} H'$. For $H' = 1 \xrightarrow{e} G'$, we obtain the *kernel* of f and the cartesian square

$$\begin{array}{ccc} \text{Ker}(f) & \longrightarrow & G \\ \downarrow & & \downarrow f \\ S & \xrightarrow{e} & G' \end{array}$$

Remark 1.12. The kernel of a homomorphism f of S -groups is for any S -scheme T given by

$$\text{Ker}(f)(T) = \ker(f(T)).$$

In particular, the $\text{Ker}(f)$ is normal.

Definition 1.13. Let G be a S -group, T a S -scheme and $g \in G(T) = \text{Hom}_S(T, G)$. Define the *lefttranslation* by g as

$$\begin{array}{ccc} G_T & \xleftarrow{=} & T \times_T G_T \\ \downarrow t_g & & \downarrow g \times \text{id} \\ G_T & \xleftarrow{m} & G_T \times_T G_T \end{array}.$$

Remark 1.14. In the situation of 1.13, for every $T' \xrightarrow{f} T$, the map

$$t_g(T'): G_T(T') = G(T') \longrightarrow G(T') = G_T(T')$$

is the lefttranslation by the element $f^*(g) \in G(T')$.

Remark 1.15. Consider

$$\begin{array}{ccc} G \times_S G & \xrightarrow{(g,h) \mapsto (gh,h)} & G \times_S G \\ \downarrow m & \nearrow \text{pr}_1 & \downarrow \\ G & & G \end{array}.$$

Let $\mathcal{P}$ be a property of morphisms stable under base change and composition with isomorphisms. Then whenever $G \rightarrow S$ satisfies $\mathcal{P}$, then m satisfies $\mathcal{P}$.